

AD-A161 132

THEORETICAL STUDIES OF LASER-INDUCED MOLECULAR RATE
PROCESSES: TOPICS IN (U) ROCHESTER UNIV NY DEPT OF
CHEMISTRY T F GEORGE OCT 85 AFOSR-TR-85-0929

1/1


UNCLASSIFIED

AFOSR-82-0046

F/G 7/4

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

OCT - 4

1

AD-A161 132

3E

MARKINGS

2

1a. REPORT SECURITY CLASSIFICATION
Unclassified

2a. SECURITY CLASSIFICATION AUTHORITY

2b. DECLASSIFICATION/DOWNGRADING SCHEDULE

3. DISTRIBUTION/AVAILABILITY OF REPORT

Approved for public release; distribution unlimited

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

5. MONITORING ORGANIZATION REPORT NUMBER(S)

AFOSR-TR 85-0929

NOV 13 1985

6a. NAME OF PERFORMING ORGANIZATION
University of Rochester6b. OFFICE SYMBOL
(If applicable)

7a. NAME OF MONITORING ORGANIZATION

AFOSR/NC

6c. ADDRESS (City, State and ZIP Code)

Department of Chemistry
River Campus
Rochester, New York 14627

7b. ADDRESS (City, State and ZIP Code)

Bldg. 410
Bolling AFB, D. C. 20332-6448

8a. NAME OF FUNDING/SPONSORING ORGANIZATION

AFOSR

8b. OFFICE SYMBOL
(If applicable)

NC

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

Grant AFOSR-82-0046

8c. ADDRESS (City, State and ZIP Code)

Bldg. 410
Bolling AFB, D. C. 20332-6448

10. SOURCE OF FUNDING NOS.

PROGRAM
ELEMENT NO.

61102F

PROJECT
NO.

2303

TASK
NO.

B3

WORK UNIT
NO.

631303

11. TITLE

THEORETICAL STUDIES OF LASER-INDUCED MOLECULAR RATE PROCESSES:
TOPICS IN LINE BROADENING AND SPECTROSCOPY

12. PERSONAL AUTHOR(S)

Thomas F. George

13a. TYPE OF REPORT

Final

13b. TIME COVERED

FROM 11/15/81 TO 9/30/85

14. DATE OF REPORT (Yr., Mo., Day)

October 1985

15. PAGE COUNT

19

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD

GROUP

SUB. GR.

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

MOLECULAR RATE PROCESSES

MOLECULAR DYNAMICS

LASER-INDUCED

LINE BROADENING

THEORETICAL STUDIES

SPECTROSCOPY

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

Theoretical studies of molecular rate processes were carried out, with an emphasis on how such processes can be induced or affected by external laser radiation. The processes were classified into two types: I Gas-Phase Processes and II Surface Processes. Within the first type a variety of laser-assisted collisional processes were investigated, including chemical reactions in a laser field, laser-induced resonance scattering with applications to transition-state spectroscopy and line broadening in general, laser-induced curve switching, isotope separation by selective radiative scattering and laser-induced associative ionization. Several related problems not necessarily involving a laser field were investigated, such as fine-structure transitions in alkali-noble-gas collisions and nonlocal potentials in bound-continuum problems. In regard to the second type of processes, the following dynamical processes were considered: electron transfer in positive-ion-surface collisions, where one-electron transfer leads to ion neutralization and two-electron transfer leads to negative-ion formation (the latter necessitates the inclusion of the Anderson correlation energy); surface-state excitation and laser-enhanced ion-semiconductor electron transfer;

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☒ DTIC USERS ☐

21. ABSTRACT SECURITY CLASSIFICATION

Unclassified

22a. NAME OF RESPONSIBLE INDIVIDUAL

Major Larry P. Davis

22b. TELEPHONE NUMBER
(Include Area Code)

(202) 767-4963

22c. OFFICE SYMBOL

NC

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

DTIC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

(19. continued) laser-induced electron-phonon processes on metal surfaces; formation of electron-hole pairs in a semiconductor by vibrationally-excited molecules; vibrational (both energy and phase) relaxation of a laser-excited adspecies; and laser-induced migration. Spectroscopic processes included resonance fluorescence of atoms near a metal surface and Stark quantum beats from beam-foil excited hydrogen atoms. Other topics of investigation were laser-induced surface excitations on a grating and static electric field effects at the surface of a metallic cluster.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

OCT - 4 1985

Final Report, October 1985

Air Force Office of Scientific Research
Grant No. AFOSR-82-0046Title: Theoretical Studies of Laser-Induced Molecular Rate
Processes: Topics in Line Broadening and Spectroscopy

Grantee: The University of Rochester

Principal Investigator: Thomas F. George
Department of Chemistry
The University of Rochester
Rochester, New York 14627Approved for public release,
distribution unlimited

85 11 12 134

ADJUTANT	
CORPORAL	<input checked="" type="checkbox"/>
DETACHMENT	<input type="checkbox"/>
UNIT	<input type="checkbox"/>

S
or.

A-1



COMPLETED PROJECT SUMMARY

1. TITLE: Theoretical Studies of Laser-Induced Molecular Rate Processes: Topics in Line Broadening and Spectroscopy
2. PRINCIPAL INVESTIGATOR: Dr. Thomas F. George
Department of Chemistry
The University of Rochester
Rochester, New York 14627
3. INCLUSIVE DATES: 15 November 1981 - 30 September 1985
4. GRANT NUMBER: AFOSR-82-0046
5. COSTS AND FY SOURCE: \$80,000, FY 82; \$100,000, FY 83;
\$107,164, FY 84; \$110,000, FY 85
6. SENIOR RESEARCH PERSONNEL:

Dr. D. Agassi	Dr. H. W. Lee
Dr. A. C. Beri	Dr. K. T. Lee
Dr. X. Y. Huang	Dr. J. T. Lin
Dr. M. Hutchinson	Dr. K. C. Liu
Dr. K. S. Lam	Mr. W. C. Murphy
Dr. I. Last	
7. JUNIOR RESEARCH PERSONNEL: Mr. F. Battaglia

8. PUBLICATIONS:

"Laser-Stimulated Molecular Dynamics and Rate Processes," T. F. George, J. Phys. Chem., 86, 10 (1982).

"On the Synergistic Effects of Laser/Phonon-Stimulated Processes: A Master Equation Approach," Surf. Sci., 115, 569 (1982).

"Recent Theoretical Chemical Dynamics at Rochester in the Paths of Joseph O. Hirschfelder," T. F. George, I. H. Zimmerman, P. L. DeVries, J. M. Yuan, K. S. Lam, D. K. Bhattacharyya and M. Hutchinson, J. Phys. Chem., 86, 1075 (1982).

"Dissociation of Adsorbed Diatomic Molecules: Combined Effects of the Surface Magnetic Field, a Laser and the Phonon Continuum," D. K. Bhattacharyya, J. Lin and T. F. George, Surf. Sci., 116, 423 (1982).

"Laser-Induced Resonance Formation in Molecular Collisions: Application of Low-Intensity Lasers to Inelastic Scattering and Spectroscopy," M. Hutchinson and T. F. George, Mol. Phys., 46, 81 (1982).

"Quantum Model of Selective Multiphoton Processes: The Role of External (Field) and Internal (Anharmonic Coupling) Detunings," J. Lin and T. F. George, Theor. Chim. Acta, 61, 243-250 (1982).

"Atomic Collisions in the Presence of Laser Radiation: Time Dependence and the Asymptotic Wavefunction," P. L. DeVries and T. F. George, Phys. Rev. A, 26, 1419-1425 (1982).

"Transition State Spectroscopy and Line Broadening," T. F. George, K. S. Lam, D. K. Bhattacharyya and M. Hutchinson, in Photon-Assisted Collisions and Related Topics, N. K. Rahman and C. Guidotti, Eds., (Harwood Academic, Chur, Switzerland, 1982), pp. 109-127.

"Laser-Stimulated Adsorption Interaction with a Semiconductor Surface," W. C. Murphy and T. F. George, J. Phys. Chem., 86, 4481 (1982).

"Quantum Model of Laser Excitation of Isotopic Species Adsorbed on a Solid Surface," J. Lin, X. Y. Huang and T. F. George, Z. Phys. B, 48, 355 (1982).

"Neutralization of Ions at an Electronically-Excited Semiconductor Surface," H. W. Lee, W. C. Murphy and T. F. George, Chem. Phys. Lett., 93, 221 (1982).

"A Semiclassical Study of Laser-Induced Atomic Fluorescence from Na_2 , K_2 and NaK ," J. M. Yuan, D. K. Bhattacharyya and T. F. George, Mol. Phys., 47, 1195 (1982).

"Laser-Generated Electron Emission from Surfaces: Effect of the Pulse Shape on Temperature and Transient Phenomena," J. Lin and T. F. George, J. Appl. Phys., 54, 382 (1983).

"Inclusion of Discrete-to-Continuum Coupling in Multiphoton Excitation and Dissociation Calculations," J. Lin and T. F. George, Theor. Chim. Acta, 62, 257 (1983).

"Semiclassical Theory of Collisional Ionization," K. S. Lam, D. K. Bhattacharyya and T. F. George, Phys. Rev. A, 27, 1353 (1983).

"Theory of Laser-Stimulated Surface Processes. I. General Formulation for the Multiphonon Relaxation of a Vibrationally Excited Adatom," A. C. Beri and T. F. George, J. Chem. Phys., 78, 4288 (1983).

"Model for the Propagation of Pulsed Surface Polaritons with Quasi-Self-Induced Transparency," X. Y. Huang, J. Lin and T. F. George, Z. Phys. B, 50, 181 (1983).

"Zeeman Transitions in Collisions of Na with Xe," T. F. George and P. L. DeVries, Chem. Phys. Lett., 96, 99 (1983).

"Quantum Dynamical Model of Laser-Stimulated Isotope Separation of Adsorbed Species: Role of Anharmonicity, Coupling Strength and Energy Feedback from the Heated Substrate," J. Lin and T. F. George, J. Chem. Phys., 78, 5197 (1983).

"Quantal Study of Laser-Induced Transitions between Electronic Potential Energy Surfaces in Reactive $\text{F} + \text{H}_2$ Collisions," I. H. Zimmerman, M. Baer and T. F. George, J. Chem. Phys., 87, 1478 (1983).

"Analysis of Laser-Enhanced Adsorption/Desorption Processes on Semiconductor Surfaces via Electronic Surface State Excitation," W. C. Murphy, A. C. Beri, T. F. George and J. Lin, in Laser Diagnostics and Photochemical Processing for Semiconductor Devices, R. M. Osgood, Jr., S. R. J. Brueck and H. R. Schlossberg, Eds., (Elsevier, New York), Mat. Res. Soc. Symp. Proc. 17, 273 (1983).

"Reply to Comments on 'Laser Excitation of Surface Electronic States for a One-Dimensional Semiconductor' by G. W. Bryant," W. C. Murphy, K. T. Lee and T. F. George, Surf. Sci., 127, L156 (1983).

"Multiphoton Resonances in Molecular Collisions," M. Hutchinson and T. F. George, J. Phys. Chem., 87, 2037 (1983).

"Quantum Statistical Theory of Laser-Stimulated Surface Processes. II. Nonequilibrium Rate of Laser Excitation and Multiphonon Relaxation," J. Lin and T. F. George, Phys. Rev. B, 28, 78 (1983).

"Quantum Dynamical Model for Laser Excitation of a Two-Level Adatom: Surface-Dressed Bloch Equations," J. Lin, X. Y. Huang and T. F. George, Solid State Commun., 47, 63 (1983).

"On the Possibility of Isotope Separation by Selective Radiative Scattering," M. Hutchinson, T. F. George and P. L. DeVries, Phys. Rev. A, 28, 490 (1983).

"Collisional Ionization as a Nonlocalized Process and the Breakdown of the Frank-Condon Approximation," K. S. Lam and T. F. George, J. Phys. Chem., 87, 2799 (1983).

"Theory of Collision-Induced Ionization of Adsorbed Species on Solid Surfaces in the Presence of Laser Radiation," K. S. Lam and T. F. George, Z. Phys. B, 51, 333 (1983).

"Theoretical Studies of Reactions in a Laser Field: $F(^2P_{3/2}, ^2P_{1/2}) + H_2 + h\nu(0.469 \text{ eV})$," I. Last, M. Baer, I. H. Zimmerman and T. F. George, Chem. Phys. Lett., 101, 163 (1983).

"Laser-Induced Gas-Phase and Surface Rate Processes: Electronic Excitations and Bound-Continuum Interactions," T. F. George, K. S. Lam, M. Hutchinson and W. C. Murphy, in Advances in Laser Spectroscopy, Vol. 2, B. A. Garetz and J. R. Lombardi, Eds., (Wiley, Chichester, England, 1983), pp. 225-242.

"Localized Magnetization Arising from Spin-Dependent Potentials in the Wolff Model," K. C. Liu, K. S. Lam and T. F. George, Phys. Status Solidi B, 120, 695 (1983).

"Emission of Short-Wavelength Photons from Ion-Surface Charge Exchange," H. W. Lee and T. F. George, IEEE J. Quantum Electron., QE-19, 1804 (1983).

"Molecular Rate Processes and Their Interaction with Laser Radiation," J. M. Yuan, P. L. DeVries, K. S. Lam and T. F. George, Prog. Phys. (Nanjing), 3, 417 (1983).

"Resonance Fluorescence of a Two-Level Atom Near a Metal Surface," X. Y. Huang, J. Lin and T. F. George, J. Chem. Phys., 80, 893 (1984).

"Nonlocal Potentials in Penning and Associative Ionization," K. S. Lam and T. G. George, Phys. Rev. A, 29, 492 (1984).

"Laser-Induced Surface Migration via Surface Plasmons," W. C. Murphy, X. Y. Huang and T. F. George, Chem. Phys. Lett., 104, 303 (1984).

"Laser-Induced Bound States at Surfaces: Ion-Neutralization and Adsorption," K. S. Lam, M. Hutchinson and T. F. George, in Collisions and Half-Collisions with Lasers, N. K. Rahman and C. Guidotti, Eds., (Harwood Academic, Chur, Switzerland, 1984), pp. 411-420.

"Laser Photochemistry," T. F. George, A. C. Beri, K. S. Lam and J. Lin, in Laser Applications, Vol. 5, J. F. Ready and R. K. Erf, Eds., (Academic Press, New York, 1984), pp. 69-127.

"Negative-Ion Formation from Surface Scattering and the Anderson Correlation Energy U," K. S. Lam, K. C. Liu and T. F. George, Phys. Lett., 101A, 356 (1984).

"Application of the Density Matrix Multiple Expansion to Stark Quantum Beats from Beam-Foil-Excited Hydrogen Atoms," F. Battaglia and T. F. George, Nuovo Cimento D, 3, 653 (1984).

"Analytic Solutions to Two-State Collision Problems for the Case of Exponential Coupling," H. W. Lee and T. F. George, Phys. Rev. A, 29, 2509 (1984).

"Laser-Induced Electron-Phonon Processes on Metal Surfaces," W. C. Murphy and T. F. George, J. Chem. Phys., 80, 5303 (1984).

"Conductivity in Semiconductors Induced by Vibrational-to-Electronic Energy Transfer," I. Last, T. F. George and D. S. Perry, Mater. Lett., 2, 315 (1984).

"Laser-Induced Neutralization and Negative-Ion Formation in Surface Scattering," K. S. Lam, K. C. Liu and T. F. George, in Laser-Assisted Deposition, Etching and Doping, S. D. Allen, Ed., Proc. Soc. Photo-Opt. Instrum. Eng. 459, 18 (1984).

"Resonance Fluorescence of Many Interacting Atoms at a Metal Surface," X. Y. Huang, T. F. George and J. Lin, in Coherence and Quantum Optics V, L. Mandel and E. Wolf, Eds. (Plenum Press, New York, 1984), pp. 685-693.

"Production of Short Wavelength (XUV) Photons from Ion-Laser-Excited-Surface Charge Exchange: Li^{3+} , He^+ + Si(111) Systems," H. W. Lee and T. F. George, in Coherence and Quantum Optics V, L. Mandel and E. Wolf, Eds., (Plenum Press, New York, 1984) pp. 923-926.

"Theory of Laser-Induced Surface Chemistry and Applications to Microelectronics and Heterogeneous Catalysis," J. Lin, W. C. Murphy and T. F. George, Ind. Eng. Chem. Prod. Res. Dev., **23**, 334 (1984).

"Resonance Fluorescence of a Two-Level Atom Near a Metal Surface. 2. Case of a Strong Driving Field," X. Y. Huang and T. F. George, J. Phys. Chem., **88**, 4801 (1984).

"Resonance Fluorescence and Photon Trapping of Two Atoms on a Metallic Surface," X. Y. Huang, K. C. Liu and T. F. George, in Laser-Controlled Chemical Processing of Surfaces, A. W. Johnson, D. J. Ehrlich and H. R. Schlossberg, Eds., (Elsevier, New York), Mat. Res. Soc. Symp. Proc., **29**, 381 (1984).

"Theoretical Aspects of Laser-Induced Periodic Surface Structure Formation," M. Hutchinson, K. T. Lee, W. C. Murphy, A. C. Beri and T. F. George, Laser-Controlled Chemical Processing of Surfaces, A. W. Johnson, D. J. Ehrlich and H. R. Schlossberg, Eds., (Elsevier, New York), Mat. Res. Soc. Symp. Proc., **29**, 389 (1984).

"How Lasers May Open the 'Last Frontier of Reaction Dynamics'," T. F. George, Ind. Chem. News, **5**, 23 (1984).

"Phase-Dissipative Mechanisms for Laser-Induced Surface Desorption/Dissociation Processes," X. Y. Huang, T. F. George, J. M. Yuan and L. M. Narducci, J. Phys. Chem., **88**, 5772 (1984).

"A Rule for the Number of Topologically Distinct Feynman Diagrams," F. Battaglia and T. F. George, J. Math. Phys., **25**, 3489 (1984); **26**, 882 (1985).

"Theory of Laser-Stimulated Surface Processes," T. F. George, J. Lin, A. C. Beri and W. C. Murphy, Prog. Surf. Sci., **16**, 139 (1984).

"Laser-Induced Molecular Dynamics: Rate Processes in the Gas Phase and at Solid Surfaces," J. Lin, M. Hutchinson and T. F. George, in Advances in Multi-Photon Processes and Spectroscopy, S. H. Lin, Ed., (World Scientific, Singapore, 1984), pp. 105-237.

"The Role of Phonons in the Excitation and Relaxation of Adspecies," A. C. Beri, K. T. Lee and T. F. George, in Proceedings of the International Conference on Lasers '83, R. C. Powell, Ed. (STS Press, McLean, Virginia, 1985), pp. 362-9.

"Negative-Ion Formation from Surface Scattering at Finite Temperatures," K. C. Liu, T. F. George and K. S. Lam, Solid State Commun., **53**, 67 (1985).

"Theoretical Studies of Chemical Reactions in a Laser Field," Sr. K. Duffy, M. Hutchinson, J. M. Yuan and T. F. George, J. Phys. (Paris), Colloq. C1, 46, (1985).

"Formation of Electron-Hole Pairs in a Semiconductor by Vibrationally-Excited Molecules," I. Last and T. F. George, Langmuir, 1, 33 (1985).

"Theoretical Study of Laser-Induced Surface Excitations on a Grating," K. T. Lee and T. F. George, Phys. Rev. B, 31, 5106 (1985).

"A Diagrammatic Approach to Ion Neutralization at Surfaces: On the Validity of First-Order Perturbation Calculations," F. Battaglia and T. F. George, J. Chem. Phys., 82, 3847 (1985).

"Recent Advances in the Theory of Chemi-Ionization," H. P. Saha, K. S. Lam and T. F. George, in Gas-Phase Chemiluminescence and Chemi-Ionization, A. Fontijn, Ed., (North Holland, Amsterdam, 1985), pp. 87-104.

"Theory of Excitation Transfer Between Rydberg Atoms," H. P. Saha and T. F. George, Mol. Phys., 55 173 (1985).

"Quantum Statistical Theory of Vibrational Dynamics in a Laser-Driven Admolecule-Surface System," X. Y. Huang, T. F. George and J. M. Yuan, J. Opt. Soc. Am. B, 2, 985 (1985).

"Static Electric Field Effects at the Surface of a Metallic Cluster," M. Hutchinson and T. F. George, Chem. Phys. Lett., 117, 468 (1985).

"Neutralization and Excitation in Low-Energy Ion-Surface Collisions," H. W. Lee and T. F. George, Surf. Sci., 159, 214 (1985).

"Multiple Resonances in Non-Franck-Condon Transitions due to Nonlocal Effects in Associative Ionization," K. S. Lam and T. F. George, Phys. Rev. A, 32, 1650 (1985).

"Theory of Laser-Stimulated Surface Processes. II. A Generalized-Master Equation Approach to Energy Transfer between an IR Laser, an Adspecies and Phonons," A. C. Beri and T. F. George, J. Chem. Phys., 83, 2482 (1985).

"Spontaneous Emission by Two Atoms with Two Different Resonance Frequencies Near a Metal Surface," K. C. Liu and T. F. George, Phys. Rev. B, 32, 3622 (1985).

"Laser-Induced Collision Processes: Polarization Effects and Ionization," K. S. Lam, P. L. DeVries and T. F. George, in Spectral Line Shapes, Vol 3, F. Rostas, Ed., (Walter de Gruyter, Berlin), in press.

"Theory of Reactions at a Solid Surface," T. F. George, K. T. Lee, W. C. Murphy, M. Hutchinson and H. W. Lee, in Theory of Chemical Reaction Dynamics, M. Baer, Ed., (CRC Press, Boca Raton, Florida), in press.

"Laser-Stimulated Vibrational Excitation of an Adspecies Studied by a Generalized Master Equation: Neutral Atomic Hydrogen on Hydrated KCl(001)," A. C. Beri and T. F. George, J. Vac. Sci. Technol. B, in press.

"Quantum Model of Dephasing-Enhanced Laser Desorption: Master Equation Approach," J. Lin, X. Y. Huang and T. F. George, J. Vac. Sci. Technol. B, in press.

"Memory Effects in Dynamical Many-Body Systems: The Isomnesic (Constant-Memory) Approximation," A. C. Beri and T. F. George, Z. Phys. B, in press.

"Perturbative Treatment of Quasi-Resonant Proton Neutralization at Alkali-Halide Surfaces," F. Battaglia, T. F. George and A. Lanaro, Surf. Sci., in press.

"Overlap Integrals for Atom-Metal Surface Interactions," W. C. Murphy and T. F. George, Int. J. Quantum Chem., in press.

"Cooperative Emission by Two Different Atoms into Surface Plasmons," K. C. Liu and T. F. George, Surf. Sci., in press.

9. ABSTRACT OF OBJECTIVES AND ACCOMPLISHMENTS:

The objectives were to help establish a theoretical foundation for the description of laser-induced molecular rate processes, including spectroscopy as well as dynamics. Two general types of processes were considered -- I. Gas-Phase Processes and II. Surface Processes -- and selected accomplishments with respect to these two general types are summarized below.

A. Gas-Phase Processes: Molecular Collisions in a Laser Field


Chemical Reactions. Quantum mechanical and semiclassical calculations were carried out for the collinear reaction $F^*(^2P_{1/2}) + H_2(v=0) - h\nu \rightarrow HF(v') + H$, where $h\nu = 0.469$ eV is the field quantum of the HF laser. It is resonant with the $v' = 1$ to $v' = 2$ vibrational transition of HF, but this resonance is not a factor in the collision and was thus ignored. Within the collision region the photon comes into resonance with two potential energy surfaces, one correlating to $F(^2P_{3/2}) + H_2$ and the other to $F^*(^2P_{1/2}) + H_2(v)$. Transmission to $HF(v') + H$ or inelastic reflection to $F(^2P_{3/2}) + H_2(v)$ takes place by stimulated emission of a photon (corresponding to the minus sign in front of $h\nu$ in the above reaction). The nominal threshold for the reactive transition to $v' = 2$ occurs at an initial translational energy of 0.004 eV. This long delay is due to the introduction of a potential barrier which is absent in the field-free case.

Curve Switching and Transition-State Spectroscopy. Laser-induced

curve switching has been studied for the process $F(^2P_{3/2}) + Xe \rightarrow F^*(^2P_{1/2}) + Xe$ in the presence of two lasers, L1 (XeCl laser) and L2 (CO_2 laser). While there is no net photon absorption from L1, the final collision energy of the $Xe + F$ system is increased (relative to the initial energy) by the energy of the photon(s) absorbed from L2. A single L2 photon is near resonant with the $v = 12$ and $v = 16$ levels of XeF , and due to the $\Delta v = 4$ transition the intensity of L2 is on the order of a MW/cm^2 , whereas L1 is only on the order of a kW/cm^2 . The initial kinetic energy was chosen low enough such that transition between the spin-orbit states of fluorine could not occur due to the collision or the "collision + L1" alone; the energy required for the spin-orbit transition was supplied by L2. An exact dynamical theory, based on a Feshbach projection operator formalism, has been derived for this process. The theory is capable of describing transition-state spectroscopy, where either L1 or L2 can be viewed as a spectroscopic probe depending on the laser frequencies and powers. Introducing suitable approximations, such as replacing certain angular momentum coupling coefficients by average values, we have carried out calculations for the kinetic-energy-averaged scattering cross section of the spin-orbit transition as a function of the frequency of L2, tuned over a range of 0.1 cm^{-1} (centered at 1076.35 cm^{-1}). For a specific rotational transition $\Delta j = 1$ of the XeF complex, the cross section forms a symmetric peak with values as high as 65 \AA^2 . We have also considered the situation with L2 turned off, where now the initial kinetic energy is greater than the fluorine

spin-orbit splitting. Plotting the cross section as a function of the frequency of L1 over a range of 0.1 cm^{-1} , we observe an asymmetric peak, where the asymmetry is due to the interference between the laser-induced resonance scattering and nonradiative potential scattering.

Isotype Separation. Related to the above work, an experimental procedure for isotope separation by selective radiative scattering has been suggested. Two colliding partners are prepared in a single beam which is formed by supersonic expansion. At some point downstream of the source, the kinetic energy of relative motion reaches a certain low value. Here two plane-polarized lasers intersect the molecular beam at right angles. Certain molecular pairs, which have the correct vibrational-rotational spacing to undergo a bound-to-bound transition $(v, l \rightarrow v', l')$, can pick up extra translational energy equal to the photon energy $h\nu_2$ of the second laser. While there is photon absorption from this laser, there is no net absorption from the first laser:



The minus sign in front of $h\nu_1$ indicates stimulated emission. Since the extra translational energy of $h\nu_2$ is much greater than the initial kinetic energy, and comparable with the forward translational energy of the beam, such molecular pairs will tend to be scattered out of the beam and can then be skimmed off. The dynamics of this process is the same as for the work described in the previous paragraph. We

have tested this hypothesis for isotope separation by using the distorted-wave Born approximation to calculate cross sections for Xe + F collisions in the presence of a UV laser ($h\nu_1$) and IR laser ($h\nu_2$). Let us define a separation efficiency p by $p = \sigma_{130} / (\sigma_{129} + \sigma_{130})$, where σ_{129} and σ_{130} are kinetic-averaged cross sections for ^{129}Xe and ^{130}Xe collisions, respectively. For laser intensities on the order of a kW/cm^2 , p is seen to be as high as 0.996. While certain aspects not taken into account in our calculations might reduce the efficiency, such as finite laser bandwidths and possible elastic collisions experienced by the deflected molecules before leaving the beam, it is clear that these velocity-changing radiative collisions should provide an efficient means of isotope separation.

B. Surface Processes

Electron Transfer in Ion-Surface Collisions. Based on the time-dependent Fano-Anderson potential, a diagrammatic expansion has been made for the reduced density matrix elements of the neutralized atom formed by one-electron transfer to an impinging positive ion. Neutralization probabilities have been computed for protons colliding with various alkali-halide surfaces, and the results are seen to be interpreted simply in terms of the collision energy and the properties of the systems, namely, the valence bandwidth of the solid and its position relative to the discrete atomic level. The theory of negative-ion formation has also been considered, where the treatment

of this two-electron transfer process now includes the effects of the Anderson correlation energy arising from the Coulombic repulsion between the two electrons of opposite spin in the same valence level of the ion. For the case of zero surface temperature, $T = 0$ K, the theory is in good agreement with molecular beam experiments on the conversion of $H^+(D^+)$ to $H^-(D^-)$ by scattering from a cesiated $W(110)$ surface. At finite temperatures, the electrons in the solid lying within the range of $k_B T$ below the Fermi level can be activated, where k_B is Boltzmann's constant. Computational results show that the thermal energy may then be used to overcome the extra energy barrier U for the second electron, provided $k_B T \geq U$, and thus enhance negative-ion formation.

Surface-State Excitation. Quantum mechanical calculations have been carried out for laser-induced excitation of electrons from the valence band to surface electronic states in a one-dimensional semiconductor (silicon). These states are localized near the surface, and excitation cross sections as high as 20 \AA^2 are obtained for low laser intensities (less than 10 W/cm^2). The calculations have been extended to include a charged adspecies. The adspecies-surface potential has been determined as a function of both the adspecies-surface separation and the laser frequency, and the surface-state excitation is seen to enhance the repulsive (attractive) part of the potential for a negative (positive) adspecies, especially as the photon energy approaches half the band-gap energy.

This idea of using a laser to "charge" the surface region has formed the basis of a semiclassical theory of charge transfer, where a positively-charged particle picks up an electron from a semiconductor surface. The charge-transfer rate is larger by a factor of $2\kappa L$ for a surface electron than for a bulk electron, where κ is the imaginary portion of the crystal momentum for the surface state and L is the length of the solid. The theory has been used to determine the probabilities of electron transfer to Li^{3+} , He^{2+} and He^+ undergoing thermal collisions with a $\text{Si}(111)$ surface. The results show an enhancement of the production of excited states of Li^{2+} , He^+ and He , respectively, receding from the surface, as compared to charge transfer in the absence of surface-state excitation. For example, for the process $\text{He}^{2+} + \text{Si}(111) + h\nu \rightarrow \text{He}^+(n=3) + \text{Si}^+$ under thermal conditions, we obtain a cross section as high as 4.19 \AA^2 , where the power density of the laser pumping the surface states is only 1 W/cm^2 (the laser photon energy $h\nu$ is chosen to be $2/5$ of the band-gap energy).

Formation of Electron-Hole Pairs in a Semiconductor by Vibrationally-Excited Molecules. A three-dimensional model of a vibrationally-excited molecule with a semiconductor has been developed, where the transfer of vibrational energy leads to the formation of electron-hole pairs.^{29,30} Results for the $\text{HCl} + \text{InSb}$ and $\text{HCl} + \text{PbSe}$ systems under thermal conditions indicate transition probabilities in the range 0.1 to 10%, which can be readily observed by experimental measurements of the semiconductor electrical conductivity. The semiconductors considered here have similar lattice

parameters. However, the transition probabilities are higher for HCl + InSb, than for HCl + PbSe, which is attributable to differences in the dielectric constant (18 for InSb and 280 for PbSe) and the energy gap (0.165 eV for InSb and 0.26 eV for PbSe -- the latter is closer to the vibrational energy level spacing of 0.358 eV in HCl).

Vibrational Energy Relaxation of a Laser-Excited Adspecies.

Vibrational energy transfer between an infrared laser, a solid surface and an adatom has been studied by a generalized master equation which includes memory effects. Numerical results using the isomnesic (constant-memory) approximation for time scales ranging from 2 picoseconds to 2 microseconds for $\text{H}(\text{H}_2\text{O})/\text{KCl}(001)$ show that bond-selective excitation is possible with a low-power laser in spite of fast phonon relaxation, and that the time evolution of the system is highly non-Markovian even for fairly long times. An overall residence time of the adspecies, up to the point of desorption, has been defined. This time is on the order of 10 nanoseconds for a laser whose power is 10 W/cm^2 and whose frequency is resonant with $v = 3$ and $v = 6$ of the adbond, where $v = 6$ is the highest vibrational level and the Debye frequency of the substrate is resonant with $v = 0$ and $v = 3$.

Laser-induced Migration. When a laser excites a plasmon on a metal surface, a wave of alternating charge is propagated along the surface. This charge has a varying electric field associated with it, and we have shown that this field can be coupled to a charged adspecies. In particular, it has been shown that if aluminum is exposed to CO_2 laser radiation of low intensity (1 W/cm^2), such a

plasmon will be excited. If an atomic oxygen ion of charge -1 were adsorbed on the surface, it would be coupled to the electric field and thus carried along with the plasmon. The velocity associated with this stimulated migration is 61.3 microns/s, which is five orders of magnitude greater than the usual thermal velocities observed at room temperature. Furthermore, the negatively-charged adspecies will only move in the direction of the plasmon; this, in turn, can be controlled by the exciting radiation. Thus a laser not only increases the speed of migration but can also select the direction.

Resonance Fluorescence at a Metal Surface. Surface-dressed optical Bloch equations, involving a treatment of surface-reflected photons and a surface plasmon, have been derived for a collision-damped two-level atom near or adsorbed on a metal surface. Effects of the laser bandwidth are included by means of a phase diffusion model for the driving field. In the weak-field or large-detuning limit, the power spectrum of scattered light has two peaks: one corresponding to Rayleigh scattering at the laser frequency ν_{21} . For a sufficiently strong driving laser field, the spectrum exhibits three peaks: a central one at $\nu_0 = \nu_L$ (Rayleigh component), a left one at $\nu_- = 2\nu_L - \nu_{21} - \delta$ (three-photon component) and a right one at $\nu_+ = \nu_{21} + \delta$ (fluorescence component), where δ is the ac Stark shift. The key feature of numerical results, which is a unique behavior due to the surface, is that for certain atom-surface distances the height H_- of the left peak is larger than H_+ of the right peak, whereas in the pure gas-phase resonance fluorescence spectrum H_- is always

less than H_+ (due to molecular collisions for the case of positive detuning).

Static Electric Field Effects at the Surface of a Metallic Cluster. A jellium/free-electron model has been used to estimate the static electric field at the surface of a spherical metallic cluster. By means of a WKB formula derived from the model, an accurate prediction can be made of the "magic numbers" (2, 10, 18, 20, 34, 40, 58, 90, 92, etc.) which refer to the enhanced abundances of certain clusters observed experimentally. Therefore, these numbers are likely to be determined by the symmetry of the problem and not by the detailed nature of the one-electron potential. The electric field near a cluster with a radius of $10\text{-}20 \text{ \AA}^2$ is found to deviate from the field at the same distance from an infinite metallic plane. Such a deviation might manifest itself by influencing physisorption.

END

FILMED

12-85

DTIC